

Chapter 17. Trigonometric Ratios

TRIGONOMETRIC RATIOS

EXERCISE - 16

1

Ⓐ Sol:

By Pythagoras theorem

$$OP^2 = OM^2 + PM^2$$

$$15^2 = 12^2 + PM^2$$


$$225 = 144 + PM^2$$

$$PM^2 = 225 - 144$$

$$PM^2 = 81$$

$$PM = \sqrt{81}$$

$$PM = 9$$



Ⓐ Sol: $\sin \theta = \frac{PM}{OP}$

$$= \frac{9}{15}$$

$$\sin \theta = \frac{3}{5}$$

Ⓑ

$$\cos \theta = \frac{OM}{OP}$$

$$= \frac{12}{15}$$

$$\cos \theta = \frac{4}{5}$$

Ⓒ

$$\tan \theta = \frac{PM}{OM}$$

$$= \frac{9}{12}$$

$$\tan \theta = \frac{3}{4}$$

$$\text{(iv)} \quad \cot \theta = \frac{OM}{PM}$$

$$= \frac{12}{9}$$

$$\cot \theta = \frac{4}{3}$$

$$\text{(v)} \quad \sec \theta = \frac{OP}{OM}$$

$$= \frac{15}{12}$$

$$\sec \theta = \frac{5}{4}$$

$$\text{(vi)} \quad \operatorname{cosec} \theta = \frac{OP}{PM}$$

$$= \frac{15}{9}$$

$$\operatorname{cosec} \theta = \frac{5}{3}$$

⑥
~~Sol~~

By Pythagoras theorem

$$AB^2 = AC^2 + BC^2$$


$$= 12^2 + 5^2$$

$$= 144 + 25$$

$$AB^2 = 169$$

$$AB = \sqrt{169}$$

$$AB = 13$$



$$\textcircled{i} \quad \sin A = \frac{BC}{AB}$$

$$\sin A = \frac{5}{13}$$

$$\textcircled{ii} \quad \cos A = \frac{AC}{AB}$$

$$\cos A = \frac{12}{13}$$

$$\textcircled{iii} \quad \sin^2 A + \cos^2 A = \left(\frac{5}{13}\right)^2 + \left(\frac{12}{13}\right)^2$$

$$= \frac{25}{169} + \frac{144}{169}$$

$$= \frac{25 + 144}{169}$$

$$= \frac{169}{169}$$

$$= 1$$

$$\therefore \sin^2 A + \cos^2 A = 1$$

$$\textcircled{iv} \quad \sec^2 A - \tan^2 A$$

$$\therefore \sec A = \frac{1}{\cos A}$$

$$= \frac{1}{\frac{12}{13}}$$

$$\sec A = \frac{13}{12}$$

$$\therefore \tan A = \frac{\sin A}{\cos A}$$

$$= \frac{\left(\frac{5}{13}\right)}{\left(\frac{12}{13}\right)}$$

$$\tan A = \frac{5}{12}$$

$$\therefore \sec^2 A - \tan^2 A = \left(\frac{13}{12}\right)^2 - \left(\frac{5}{12}\right)^2$$

$$= \frac{169}{144} - \frac{25}{144}$$

$$= \frac{169-25}{144}$$

$$= \frac{144}{144}$$

$$= 1.$$

$$\therefore \sec^2 A - \tan^2 A = 1.$$

2
sol.

(a)

By pythagorus theorem

\therefore hypotenuse = BC

$$\therefore BC^2 = AB^2 + AC^2$$

$$10^2 = 6^2 + AC^2$$


$$100 = 36 + AC^2$$

$$AC^2 = 100 - 36$$

$$AC^2 = 64$$

$$AC = \sqrt{64}$$

$$AC = 8.$$



$$\textcircled{i} \quad \sin B = \frac{Ac}{Bc}$$

$$= \frac{8}{10}$$

$$\sin B = \frac{4}{5}$$

$$\textcircled{ii} \quad \cos c = \frac{Ac}{Bc}$$

$$= \frac{8}{10}$$

$$\cos c = \frac{4}{5}$$

$$\textcircled{iii} \quad \sin B + \sin c$$

$$\therefore \sin c = \frac{AB}{Bc}$$

$$= \frac{6}{10}$$

$$= \frac{3}{5}$$

$$\therefore \sin B = \frac{4}{5}$$

$$\therefore \sin B + \sin c = \frac{4}{5} + \frac{3}{5}$$

$$= \frac{4+3}{5}$$

$$\sin B + \sin c = \frac{7}{5}$$

$$\textcircled{iv} \quad \sin B \cdot \cos c + \sin c \cdot \cos B$$

$$\cos B = \frac{AB}{Bc} = \frac{6}{10}$$

$$= \frac{3}{5}$$

$$\therefore \sin B \cdot \cos C + \sin C \cdot \cos B$$

$$\Rightarrow \frac{4}{5} \cdot \frac{4}{5} + \frac{3}{4} \cdot \frac{3}{5}$$

$$\Rightarrow \frac{16}{25} + \frac{9}{25}$$

$$\Rightarrow \frac{16+9}{25}$$

$$\Rightarrow \frac{25}{25}$$

$$\Rightarrow 1$$

⑥

From Figure

$\triangle ADC$

$$\Rightarrow AD^2 + CD^2 = AC^2$$

$$AD^2 + 5^2 = 13^2$$


$$AD^2 + 25 = 169$$

$$AD^2 = 169 - 25$$

$$AD^2 = 144$$

$$AD = \sqrt{144}$$

$$AD = 12$$



From Figure

$\triangle ABD$

$$\Rightarrow BD = BC - CD$$

$$= 21 - 5$$

$$BD = 16$$

$$\therefore AB^2 = BD^2 + AD^2$$

$$AB^2 = 16^2 + 12^2$$

$$AB^2 = 256 + 144$$

$$AB^2 = 400$$

$$AB = \sqrt{400}$$

$$AB = 20$$

$$\therefore AB = 20, AD = 12;$$

$$BD = 16$$

$$AC = 13$$

$$CD = 5$$

$$\textcircled{i} \quad \tan x = \frac{CD}{AD}$$

$$\tan x = \frac{5}{12}$$

$$\textcircled{ii} \quad \cos y = \frac{BD}{AB}$$

$$= \frac{16}{20}$$

$$\cos y = \frac{4}{5}$$

$$\textcircled{iii} \quad \operatorname{cosec}^2 y - \cot^2 y$$

$$\rightarrow \operatorname{cosec} y = \frac{AB}{AD}$$

$$= \frac{20}{12}$$

$$\operatorname{cosec} y = \frac{5}{3}$$

$$\Rightarrow \cot y = \frac{BD}{AD}$$

$$= \frac{16}{12}$$

$$\cot y = \frac{4}{3}$$

$$\therefore \operatorname{cosec}^2 y - \cot^2 y = \left(\frac{5}{3}\right)^2 - \left(\frac{4}{3}\right)^2$$

$$= \frac{25}{9} - \frac{16}{9}$$

$$= \frac{25-16}{9}$$

$$= \frac{9}{9}$$

$$\therefore \operatorname{cosec}^2 y - \cot^2 y = 1$$

③

sol:

①

From figure

$$BD = BC - CD$$

$$BD = 21 - 5$$

$$BD = 16$$

From $\triangle ADC$

$$AD^2 + DC^2 = AC^2$$

$$AD^2 + 5^2 = 13^2$$


$$AD^2 + 25 = 169$$

$$AD^2 = 169 - 25$$

$$AD^2 = 144$$

$$AD = \sqrt{144}$$

$$AD = 12$$



$$\therefore \sec \theta = \frac{AB}{BD}$$

from fig $\triangle ABD$

$$\begin{aligned} AB^2 &= BD^2 + AD^2 \\ &= 16^2 + 12^2 \\ &= 256 + 144 \\ &= 400 \end{aligned}$$

$$AB^2 = 400$$

$$AB = \sqrt{400}$$

$$AB = 20$$

$$\therefore \sec \theta = \frac{20}{16}$$

$$= \frac{5}{4}$$

(6)

sol:

From figure

$\triangle ABC$

$$\begin{aligned} AC^2 &= AB^2 + BC^2 \\ &= 3^2 + 4^2 \\ &= 9 + 16 \end{aligned}$$

$$AC^2 = 25$$


$$AC = \sqrt{25}$$

$$AC = 5$$

From figure

$\triangle BCD$

$$\begin{aligned} CD^2 &= BC^2 + BD^2 \\ 12^2 &= 4^2 + BD^2 \end{aligned}$$



$$144 = 16 + BD^2$$

$$BD^2 = 144 - 16$$

$$= 128$$

$$BD^2 = 128$$

$$BD = \sqrt{128}$$

$$= \sqrt{16 \times 4 \times 2}$$

$$= 4 \times 2\sqrt{2}$$

$$BD = 8\sqrt{2}$$

∴

$$\textcircled{i} \quad \sin x = \frac{BC}{AC}$$

$$\sin x = \frac{4}{5}$$

$$\textcircled{ii} \quad \cot x = \frac{AB}{BC}$$

$$= \frac{3}{4}$$

$$\textcircled{iii} \quad \cot^2 x - \operatorname{cosec}^2 x$$

$$\therefore \cot x = \frac{3}{4}$$

$$\operatorname{cosec} x = \frac{AC}{BC}$$

$$= \frac{5}{4}$$

$$\therefore \cot^2 x - \operatorname{cosec}^2 x = \left(\frac{3}{4}\right)^2 - \left(\frac{5}{4}\right)^2$$

$$= \frac{9}{16} - \frac{25}{4}$$

$$\begin{aligned} \therefore \cot^2 x - \operatorname{cosec}^2 x &= \frac{9 - 25}{16} \\ &= \frac{-16}{16} \\ &= -1 \end{aligned}$$

$$\begin{aligned} \text{(iv) } \sec y &= \frac{CP}{BD} \\ &= \frac{12}{8\sqrt{2}} \end{aligned}$$

$$\sec y = \frac{3}{2\sqrt{2}}$$

$$\text{(v) } \tan^2 y = \frac{1}{\cos^2 y}$$

$$\begin{aligned} \therefore \sec^2 y &= \frac{1}{\cos^2 y} \\ &= \left(\frac{3}{2\sqrt{2}}\right)^2 \\ &= \frac{9}{4 \times 2} \end{aligned}$$

$$\frac{1}{\cos^2 y} = \frac{9}{8}$$

$$\begin{aligned} \therefore \tan^2 y &= \left(\frac{BC}{BD}\right)^2 \\ &= \left(\frac{4}{8\sqrt{2}}\right)^2 \\ &= \frac{16}{64 \times 2} \\ &= \frac{1}{4} \end{aligned}$$

$$\tan^2 y = \frac{1}{8}$$

$$\begin{aligned} \therefore \tan y - \frac{1}{\cos y} &= \frac{1}{8} - \frac{9}{8} \\ &= \frac{1-9}{8} \\ &= \frac{-8}{8} \\ &= -1 \end{aligned}$$

④

Sol

⑨

From figure

 $\triangle BCD$

$$\Rightarrow BC^2 = BD^2 + CD^2$$


$$BC^2 = 9^2 + 12^2$$

$$= 81 + 144$$

$$BC^2 = 225$$

$$BC = \sqrt{225}$$

$$BC = 15.$$



$$\begin{aligned} \triangle ABC \Rightarrow AB^2 + BC^2 &= AC^2 \\ AB^2 + 15^2 &= 25^2 \\ AB^2 &= 625 - 225 \\ AB^2 &= 400 \\ AB &= \sqrt{400} = 20. \end{aligned}$$

$$\textcircled{i} \quad 2 \sin y - \cos y$$

$$\therefore \sin y = \frac{BD}{BC}$$

$$= \frac{9}{15}$$

$$\therefore \sin y = \frac{3}{5}$$

$$\cos y = \frac{CD}{BC}$$

$$= \frac{12}{15}$$

$$\therefore \cos y = \frac{4}{5}$$

$$\therefore 2 \sin y - \cos y = 2 \cdot \frac{3}{5} - \frac{4}{5}$$

$$= \frac{6}{5} - \frac{4}{5}$$

$$= \frac{6-4}{5}$$

$$= \frac{2}{5} //$$

$$\textcircled{ii} \quad 2 \sin x - \cos x$$

$$\sin x = \frac{BC}{AC} = \frac{15}{25} = \frac{3}{5}$$

$$\cos x = \frac{AB}{AC} = \frac{20}{25} = \frac{4}{5}$$

$$\therefore 2 \sin x - \cos x = 2 \cdot \frac{3}{5} - \frac{4}{5}$$

$$= \frac{6}{5} - \frac{4}{5}$$

$$= \frac{6-4}{5} = \frac{2}{5}$$

$$(iii) \quad 1 - \sin x + \cos y$$

$$\therefore \sin x = \frac{3}{5}$$

$$\cos y = \frac{4}{5}$$

$$\begin{aligned} \therefore 1 - \sin x + \cos y &= 1 - \frac{3}{5} + \frac{4}{5} \\ &= \frac{5 - 3 + 4}{5} \\ &= \frac{6}{5} \end{aligned}$$

$$(iv) \quad 2 \cos x - 3 \sin y + 4 \tan x$$

$$\therefore \sin x = \frac{3}{5}$$

$$\cos x = \frac{4}{5}$$

$$\sin y = \frac{3}{5}$$

$$\begin{aligned} \therefore \tan x &= \frac{\sin x}{\cos x} \\ &= \frac{\frac{3}{5}}{\frac{4}{5}} \end{aligned}$$

$$\tan x = \frac{3}{4}$$

$$\therefore 2 \cos x - 3 \sin y + 4 \tan x$$

$$\rightarrow 2 \cdot \frac{4}{5} - 3 \cdot \frac{3}{5} + 4 \cdot \frac{3}{4}$$

$$\Rightarrow \frac{8}{5} - \frac{9}{5} + 3$$

$$\rightarrow \frac{8 - 9 + 15}{5}$$

$$\Rightarrow \frac{14}{5}$$

(b)

Sol:

(ii)

By pythagorus theorem

$$AC^2 = BC^2 + AB^2$$

$$5^2 = 3^2 + AB^2$$

$$25 = 9 + AB^2$$


$$AB^2 = 25 - 9$$

$$AB^2 = 16$$

$$AB = \sqrt{16}$$

$$AB = 4$$

$$\therefore AB = y = 4.$$




(i)

$$\sin x^\circ = \frac{BC}{AC}$$

$$\sin x^\circ = \frac{3}{5}$$

5Sol:

$$\text{Given } \tan A = \frac{yz}{xy} = \frac{5}{12}$$



By pythagorus theorem

$$xy^2 + yz^2 = xz^2$$

$$12^2 + 5^2 = xz^2$$

$$\therefore 144 + 25 = xz^2$$

$$xz^2 = 169$$

$$xz = 13$$

$$(i) \quad \cos A = \frac{x_4}{x_3}$$

$$= \frac{12}{13}$$

$$(ii) \quad \operatorname{cosec} A - \cot A$$

$$\operatorname{cosec} A = \frac{x_3}{y_2}$$

$$= \frac{13}{5}$$

$$\cot A = \frac{x_4}{y_3}$$

$$= \frac{12}{5}$$

$$\therefore \operatorname{cosec} A - \cot A = \frac{13}{5} - \frac{12}{5}$$

$$= \frac{13-12}{5}$$

$$= \frac{1}{5}$$

6

sol:

(a)

Given $AB = 7$

$$BC - AC = 1$$

By Pythagoras theorem


$$BC^2 = AB^2 + AC^2$$

$$\therefore BC = 1 + AC$$

$$\therefore (1 + AC)^2 = 7^2 + AC^2$$

$$1 + AC^2 + 2AC = 49 + AC^2$$

$$2AC = 49 - 1$$



$$2AC = 48$$

$$AC = 24$$

$$(i) \sin C = \frac{AB}{BC}$$

\therefore from given

$$BC - AC = 1$$

$$BC - 24 = 1$$

$$BC = 1 + 24$$

$$BC = 25$$

$$\sin C = \frac{7}{25}$$

$$(ii) \tan B = \frac{AC}{AB}$$

$$= \frac{24}{7}$$


7
81:

(i)

$$\cos \angle ABC = \frac{BD}{BA}$$

$$= \frac{9}{18.5}$$

$$= \frac{3}{5}$$



$$\begin{aligned}
 \text{(ii)} \quad \sin \angle ACB &= \frac{AD}{AC} \\
 &= \frac{12}{15} \\
 &= \frac{4}{5}
 \end{aligned}$$

6
6
50

$$\text{Given } PQ = 40$$

$$PR + QR = 50$$

By pythagoruous theorem

$$PR^2 = PQ^2 + QR^2$$

$$(50 - QR)^2 = PQ^2 + QR^2$$

$$2500 + QR^2 - 100QR = 40^2 + QR^2$$

$$2500 - 1600 = 100QR$$

$$100QR = 900$$

$$QR = \frac{900}{100}$$


$$QR = 9.$$

$$\text{Given } PR + QR = 50$$

$$PR + 9 = 50$$

$$PR = 50 - 9$$

$$PR = 41$$



$$\begin{aligned} \text{(i)} \quad \sin P &= \frac{QR}{PR} \\ &= \frac{9}{41} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \cos P &= \frac{PQ}{PR} \\ &= \frac{40}{41} \end{aligned}$$

$$\text{(iii)} \quad \tan R = \frac{\sin R}{\cos R}$$


$$\begin{aligned} \therefore \sin R &= \frac{PQ}{PR} \\ &= \frac{40}{41} \end{aligned}$$

$$\begin{aligned} \cos R &= \frac{QR}{PR} \\ &= \frac{9}{41} \end{aligned}$$

$$\begin{aligned} \therefore \tan R &= \frac{\frac{40}{41}}{\frac{9}{41}} \\ &= \frac{40}{9} \end{aligned}$$

$$\therefore \tan R = \frac{40}{9}$$

Q. 10



Given $AB = AC = 5 \text{ cm}$

$BC = 6$

$$(i) \quad \sin C = \frac{AD}{AC}$$

$$= \frac{4}{5}$$

$$(ii) \quad \tan B = \frac{AD}{BD}$$

$$= \frac{4}{3}$$

(iii) $\tan C - \cot B$

$$\therefore \tan C = \frac{AD}{DC} = \frac{4}{3}$$

$$\cot B = \frac{BD}{AD} = \frac{3}{4}$$

$$\therefore \tan C - \cot B = \frac{4}{3} - \frac{3}{4}$$

$$= \frac{16-9}{12}$$


$$= \frac{7}{12}$$

(b)

Given $AB = 2$
 $BC = 1$

$$\therefore \sin \theta = \frac{AB}{AC}$$

$$\tan \theta = \frac{AB}{BC}$$



\therefore By Pythagorean theorem

$$AC^2 = AB^2 + BC^2$$

$$AC^2 = 2^2 + 1^2$$

$$AC^2 = 4 + 1$$

$$AC^2 = 5$$

$$AC = \sqrt{5}$$

$$\therefore \sin^2 \theta = \left(\frac{AB}{AC}\right)^2 = \left(\frac{2}{\sqrt{5}}\right)^2 = \frac{4}{5}$$

$$\tan^2 \theta = \left(\frac{AB}{BC}\right)^2 = \left(\frac{2}{1}\right)^2 = 4$$

$$\therefore \sin^2 \theta + \tan^2 \theta = \frac{4}{5} + 4$$

$$= \frac{4+20}{5}$$

$$= \frac{24}{5}$$


$$\sin^2 \theta + \tan^2 \theta = \underline{\underline{\frac{24}{5}}}$$

(c)

Given $BD = 15$

$$\sin B = \frac{4}{5}$$

$$\tan C = 1$$



$$\therefore \sin B = \frac{4}{5} \times \frac{5}{5}$$

$$\sin B = \frac{AD}{AB} = \frac{20}{25}$$

$$(i) \therefore AD = 20$$

$$\therefore \tan C \Rightarrow \frac{\sin C}{\cos C} = 1$$

$$\Rightarrow \frac{AD}{DC} = 1$$

$$\therefore AD = DC$$

$$\therefore DC = 20$$

In $\triangle ACD$; By Pythagoras theorem

$$\begin{aligned} AC^2 &= AD^2 + DC^2 \\ &= 20^2 + 20^2 \\ &= 400 + 400 \end{aligned}$$

$$AC^2 = 800$$

$$AC = \sqrt{800}$$

$$= \sqrt{400 \times 2}$$

$$AC = 20\sqrt{2}$$

$$(ii) \tan^2 B - \frac{1}{\cos^2 B} = -1.$$

$$\therefore \text{LHS} \Rightarrow \tan^2 B - \frac{1}{\cos^2 B}$$

$$\begin{aligned} \tan^2 B &= \left(\frac{AD}{BD}\right)^2 = \left(\frac{20}{15}\right)^2 = \left(\frac{4}{3}\right)^2 \\ &= \frac{16}{9} \end{aligned}$$

$$\cos^2 B = \left(\frac{BD}{AB}\right)^2 = \left(\frac{15}{25}\right)^2 = \left(\frac{3}{5}\right)^2 = \frac{9}{25}$$

$$\therefore \tan^2 B - \frac{1}{\cos^2 B} \Rightarrow \frac{16}{9} - \frac{1}{\left(\frac{9}{25}\right)}$$

$$\Rightarrow \frac{16}{9} - \frac{25}{9}$$

$$\Rightarrow \frac{16-25}{9}$$

$$\Rightarrow -1.$$

$$\therefore \text{LHS} = \text{RHS}$$

Hence proved.

18/10
①

Given


$$\sin \theta = \frac{3}{5}$$

\therefore From Pythagoras theorem

$$b^2 = a^2 + c^2$$

$$25 = 9 + c^2$$

$$25 = 9 + c^2$$



$$ac^2 = 25 - 9$$

$$ac^2 = 16$$

$$ac = \sqrt{16}$$

$$ac = 4$$

$$\begin{aligned} \therefore \cos \theta &= \frac{ac}{bc} \\ &= \frac{4}{5} \end{aligned}$$

(ii)

$$\begin{aligned} \tan \theta &= \frac{\sin \theta}{\cos \theta} \\ &= \frac{\frac{3}{5}}{\frac{4}{5}} \\ &= \frac{3}{4} \end{aligned}$$

(10)
Sol:Given that $\tan \theta = \frac{5}{12}$


By pythagorus theorem

$$\begin{aligned} bc^2 &= ab^2 + ac^2 \\ &= 5^2 + 12^2 \\ &= 25 + 144 \end{aligned}$$

$$bc^2 = 169$$

$$bc = \sqrt{169}$$

$$\therefore bc = 13$$



$$\begin{aligned}\sin \theta &= \frac{ab}{bc} \\ &= \frac{5}{13}\end{aligned}$$

$$\begin{aligned}\cos \theta &= \frac{ac}{bc} \\ &= \frac{12}{13}\end{aligned}$$

⑪
Sol:

Given $\sin \theta = \frac{6}{10}$

by pythagorus theorem

$$bc^2 = ab^2 + ac^2$$

$$10^2 = 6^2 + ac^2$$

$$100 = 36 + ac^2$$

$$ac^2 = 100 - 36$$

$$ac^2 = 64$$

$$ac = \sqrt{64}$$


$$ac = 8$$

$$\therefore \cos \theta = \frac{ac}{bc} = \frac{8}{10}$$

$$\tan \theta = \frac{ab}{ac} = \frac{6}{8}$$


$$\therefore \cos \theta + \tan \theta = \frac{8}{10} + \frac{6}{8} = \frac{64 + 60}{80}$$

$$= \frac{124}{80} \Rightarrow \frac{31}{20} = 1\frac{11}{20}$$



(12)
Sol

Given $\tan \theta = \frac{4}{3}$



by pythagorous theorem

$$ab^2 + ac^2 = bc^2$$

$$4^2 + 3^2 = bc^2$$

$$16 + 9 = bc^2$$

$$bc^2 = 25$$

$$bc = \sqrt{25}$$

$$bc = 5.$$

$$\therefore \sin \theta = \frac{ab}{bc} = \frac{4}{5}$$

$$\cos \theta = \frac{ac}{bc} = \frac{3}{5}$$

$$\therefore \sin \theta + \cos \theta = \frac{4}{5} + \frac{3}{5}$$

$$= \frac{4+3}{5}$$


$$= \frac{7}{5}$$

(13)
Sol

Given $\operatorname{cosec} \theta = \sqrt{5}$

$$\operatorname{cosec} \theta = \frac{1}{\sin \theta} = \frac{bc}{ab} = \frac{\sqrt{5}}{1}$$

$$\therefore bc = \sqrt{5} ; ab = 1$$



By pythagorus theorem,

$$bc^2 = ab^2 + ac^2$$

$$(\sqrt{5})^2 = 1^2 + ac^2$$

27

$$5 = 1 + ac^2$$

$$ac^2 = 5 - 1$$

$$ac^2 = 4$$

$$ac = \sqrt{4}$$

$$ac = 2$$

$$\therefore \cot \theta - \cos \theta = \frac{ac}{ab} - \frac{ac}{bc}$$

$$= \frac{2}{1} - \frac{2}{\sqrt{5}}$$

$$= \frac{2\sqrt{5} - 2}{\sqrt{5}}$$

$$= \frac{2(\sqrt{5} - 1)}{\sqrt{5}}$$

Solution - 14 :-

$$\text{Given } \sin \theta = \frac{p}{q}$$

by pythagoras theorem


$$q^2 = p^2 + ac^2$$

$$ac^2 = q^2 - p^2$$

$$ac = \sqrt{q^2 - p^2}$$

$$\cos \theta = \frac{ac}{bc} = \frac{\sqrt{q^2 - p^2}}{q}$$

$$\therefore \cos \theta + \sin \theta = \frac{\sqrt{q^2 - p^2}}{q} + \frac{p}{q} = \frac{p + \sqrt{q^2 - p^2}}{q}$$



Solution-15

Given $\tan \theta = \frac{8}{15}$

by pythagorus theorem

$$bc^2 = ab^2 + ac^2$$

$$bc^2 = 8^2 + 15^2$$

$$bc^2 = 64 + 225$$

$$bc^2 = 289$$

$$bc = \sqrt{289}$$

$$bc = 17$$

$$\therefore ab=8 ; ac=15 ; bc=17$$

$$\sec \theta = \frac{bc}{ac} = \frac{17}{15}$$

$$\operatorname{cosec} \theta = \frac{bc}{ab} = \frac{17}{8}$$


$$\therefore \sec \theta + \operatorname{cosec} \theta = \frac{17}{15} + \frac{17}{8}$$

$$= \frac{17 \times 8 + 17 \times 15}{120}$$

$$= \frac{391}{120}$$

$$= 3 \frac{31}{120}$$

28



Solution-16:


Given $13 \sin A = 5$

$$\sin A = \frac{5}{13}$$

by pythagorus theorem

$$BA^2 = AB^2 + AC^2$$

$$\rightarrow 13^2 = 5^2 + AC^2$$



$$169 = 25 + AC^2$$

$$AC^2 = 169 - 25$$

$$AC^2 = 144$$

$$AC = \sqrt{144}$$

$$AC = 12$$

$$\therefore AB = 5 ; BA = 13 ; AC = 12$$

$$\sin A = \frac{5}{13}$$

$$\cos A = \frac{12}{13}$$

$$\begin{aligned} \tan A &= \frac{\sin A}{\cos A} \\ &= \frac{\left(\frac{5}{13}\right)}{\left(\frac{12}{13}\right)} \\ &\Rightarrow \frac{5}{12} \end{aligned}$$

$$\begin{aligned} \therefore \frac{5 \sin A - 2 \cos A}{\tan A} &= \frac{5 \left(\frac{5}{13}\right) - 2 \left(\frac{12}{13}\right)}{\left(\frac{5}{12}\right)} \\ &= \frac{\frac{25}{13} - \frac{24}{13}}{\frac{5}{12}} \\ &= \frac{\frac{25-24}{13}}{\frac{5}{12}} \\ &= \frac{1}{13} \times \frac{12}{5} \Rightarrow \frac{12}{65} \end{aligned}$$

Solution -17

Given $\operatorname{cosec} A = \sqrt{2}$
 $\operatorname{cosec} A = \frac{1}{\sin A} = \sqrt{2}$

$$\therefore \sin A = \frac{1}{\sqrt{2}}$$

By pythagoras theorem

$$bc^2 = ab^2 + ac^2$$

$$(\sqrt{2})^2 = 1^2 + ac^2$$

$$2 = 1 + ac^2$$

$$ac^2 = 2 - 1$$

$$ac^2 = 1$$

$$ac = \sqrt{1}$$

$$ac = 1$$

$$\therefore ac = 1 ; bc = \sqrt{2} ; ab = 1$$

$$\therefore \sin A = \frac{1}{\sqrt{2}}$$


$$\cos A = \frac{1}{\sqrt{2}}$$

$$\tan A = \frac{\sin A}{\cos A} = \frac{\left(\frac{1}{\sqrt{2}}\right)}{\left(\frac{1}{\sqrt{2}}\right)} = 1$$

$$\cot A = \frac{1}{\tan A}$$

$$= \frac{1}{1}$$

$$\cot A = 1$$



$$\therefore \frac{2 \sin^2 A + 3 \cot^2 A}{\tan^2 A - \cos^2 A}$$

$$\Rightarrow \frac{2 \cdot \left(\frac{1}{\sqrt{2}}\right)^2 + 3(1)^2}{(1)^2 - \left(\frac{1}{\sqrt{2}}\right)^2}$$

$$\Rightarrow \frac{2 \cdot \frac{1}{2} + 3 \cdot (1)}{1 - \frac{1}{2}}$$

$$\Rightarrow \frac{1+3}{\left(\frac{2-1}{2}\right)}$$

$$\Rightarrow \frac{4 \times 2}{1}$$

$$\Rightarrow 8.$$

Solution 18 :-

Given ABCD is a rhombus

$$AC = 8 ; BD = 6$$

From figure

$\triangle OBC$;



By pythagorus theorem

$$BC^2 = OC^2 + OB^2$$

$$= 3^2 + 4^2$$

$$= 9 + 16 = 25$$

$$BC = \sqrt{25} = 5$$



$$\begin{aligned}\sin \angle OCB &= \frac{OB}{BC} \\ &= \frac{3}{5}\end{aligned}$$

Solution-19:-

$$\text{Given } \tan \theta = \frac{5}{12}$$

By pythagoras theorem

$$bc^2 = ab^2 + ac^2$$

$$bc^2 = 5^2 + 12^2$$

$$bc^2 = 25 + 144$$

$$bc^2 = 169$$

$$bc = \sqrt{169}$$

$$bc = 13$$


$$\therefore bc = 13; ab = 5, ac = 12$$

$$\therefore \sin \theta = \frac{ab}{bc} = \frac{5}{13}$$

$$\cos \theta = \frac{ac}{bc} = \frac{12}{13}$$

$$\frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta} = \frac{\frac{12}{13} + \frac{5}{13}}{\frac{12}{13} - \frac{5}{13}}$$

$$= \frac{\frac{12+5}{13}}{\frac{12-5}{13}} = \frac{17}{78} = 2\frac{3}{7} //$$



Solution - 20 :

$$\text{Given } 5\cos A - 12\sin A = 0$$

$$5\cos A = 12\sin A$$

$$\frac{5}{12} = \frac{\sin A}{\cos A}$$

$$\frac{5}{12} = \tan A$$

By Pythagoras theorem

$$bc^2 = ab^2 + ac^2$$

$$bc^2 = 5^2 + 12^2$$

$$bc^2 = 25 + 144$$

$$bc^2 = 169$$

$$bc = \sqrt{169}$$

$$bc = 13$$

$$\therefore ab = 5 ; bc = 13 ; ac = 12$$


$$\sin A = \frac{ab}{bc} = \frac{5}{13}$$

$$\cos A = \frac{ac}{bc} = \frac{12}{13}$$

$$\therefore \frac{\sin A + \cos A}{2\cos A - \sin A} = \frac{\frac{5}{13} + \frac{12}{13}}{2 \cdot \frac{12}{13} - \frac{5}{13}}$$

$$= \frac{(5+12)/13}{(24-5)/13}$$

$$= \frac{17}{19}$$



Solution - 21


Given $\tan \theta = \frac{p}{q}$

by pythagorus theorem

$$bc^2 = ab^2 + ac^2$$

$$bc^2 = p^2 + q^2$$

$$bc = \sqrt{p^2 + q^2}$$



$$\therefore \sin \theta = \frac{ab}{bc} = \frac{p}{\sqrt{p^2 + q^2}}$$

$$\cos \theta = \frac{ac}{bc} = \frac{q}{\sqrt{p^2 + q^2}}$$

$$\therefore \frac{p \sin \theta - q \cos \theta}{p \sin \theta + q \cos \theta} = \frac{p \cdot \frac{p}{\sqrt{p^2 + q^2}} - q \cdot \frac{q}{\sqrt{p^2 + q^2}}}{p \cdot \frac{p}{\sqrt{p^2 + q^2}} + q \cdot \frac{q}{\sqrt{p^2 + q^2}}}$$

$$= \frac{p^2 - q^2}{\sqrt{p^2 + q^2}}$$

$$\frac{p^2 - q^2}{\sqrt{p^2 + q^2}}$$

$$= \frac{p^2 - q^2}{p^2 + q^2}$$

Solution - 22 :

35

$$\begin{aligned}\text{Given } 3 \cot \theta &= 4 \\ \cot \theta &= \frac{4}{3}\end{aligned}$$

By pythagoras theorem

$$bc^2 = ab^2 + ac^2$$

$$bc^2 = 3^2 + 4^2$$

$$bc^2 = 9 + 16$$

$$bc^2 = 25$$

$$bc = \sqrt{25}$$

$$bc = 5$$

$$\therefore \sin \theta = \frac{ab}{bc} = \frac{3}{5}$$


$$\cos \theta = \frac{ac}{bc} = \frac{4}{5}$$

$$\therefore \frac{5 \sin \theta - 3 \cos \theta}{5 \sin \theta + 3 \cos \theta} = \frac{5 \cdot \frac{3}{5} - 3 \cdot \frac{4}{5}}{5 \cdot \frac{3}{5} + 3 \cdot \frac{4}{5}}$$

$$= \frac{(15 - 12) / 5}{(15 + 12) / 5}$$

$$= \frac{3}{27}$$

$$= \frac{1}{9}$$



Solution - 23 (i)


36

Given $5\cos\theta - 12\sin\theta = 0$

$$5\cos\theta = 12\sin\theta$$

$$\frac{5}{12} = \frac{\sin\theta}{\cos\theta}$$

$$\frac{5}{12} = \tan\theta$$



By Pythagoras theorem

$$bc^2 = ab^2 + ac^2$$

$$bc^2 = 5^2 + 12^2$$

$$bc^2 = 25 + 144$$

$$bc^2 = 169$$

$$bc = \sqrt{169}$$

$$bc = 13$$

$$\therefore \sin\theta = \frac{ab}{bc} = \frac{5}{13}$$

$$\cos\theta = \frac{ac}{bc} = \frac{12}{13}$$

$$\therefore \frac{\sin\theta + \cos\theta}{2\cos\theta - \sin\theta} = \frac{\frac{5}{13} + \frac{12}{13}}{2 \cdot \frac{12}{13} - \frac{5}{13}}$$

$$= \frac{\frac{5+12}{13}}{\frac{24-5}{13}}$$

$$= \frac{17}{13} //$$


Solution-23 (ii) :-

37

Given $\operatorname{cosec} \theta = \frac{13}{12}$

$$\Rightarrow \operatorname{cosec} \theta = \frac{1}{\sin \theta} = \frac{13}{12}$$

$$\therefore \sin \theta = \frac{12}{13}$$



By pythagorus theorem

$$bc^2 = ab^2 + ac^2$$

$$13^2 = 12^2 + ac^2$$

$$169 = 144 + ac^2$$

$$169 - 144 = ac^2$$

$$25 = ac^2$$

$$ac = \sqrt{25}$$

$$ac = 5$$

$$\therefore \sin \theta = \frac{12}{13}$$

$$\cos \theta = \frac{ac}{bc} = \frac{5}{13}$$

$$\therefore \frac{2 \sin \theta - 3 \cos \theta}{4 \sin \theta - 9 \cos \theta} = \frac{2 \cdot \frac{12}{13} - 3 \cdot \frac{5}{13}}{4 \cdot \frac{12}{13} - 9 \cdot \frac{5}{13}}$$

$$= \frac{24 - 15}{48 - 45}$$

$$= \frac{9}{3} = \underline{\underline{3}}$$

Solution - 24 :-

38

Given $5 \sin \theta = 3$

$$\sin \theta = \frac{3}{5}$$

By Pythagoras theorem

$$bc^2 = ab^2 + ac^2$$

$$5^2 = 3^2 + ac^2$$

$$25 = 9 + ac^2$$

$$ac^2 = 25 - 9$$

$$ac^2 = 16$$

$$ac = \sqrt{16}$$


$$ac = 4.$$

$$\cos \theta = \frac{ac}{bc} = \frac{4}{5}$$

$$\therefore \sec \theta = \frac{1}{\cos \theta} = \frac{5}{4}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{3/5}{4/5} = \frac{3}{4}$$

$$\begin{aligned} \therefore \frac{\sec \theta - \tan \theta}{\sec \theta + \tan \theta} &= \frac{\frac{5}{4} - \frac{3}{4}}{\frac{5}{4} + \frac{3}{4}} \\ &= \frac{\frac{5-3}{4}}{\frac{5+3}{4}} \\ &= \frac{2}{8} \\ &= \frac{1}{4}. \end{aligned}$$



Solution - 25 :-

Given $\sin \theta = \cos \theta$

Then $2 \tan^2 \theta + \sin^2 \theta - 1$

$$\Rightarrow 2 \left(\frac{\sin \theta}{\cos \theta} \right)^2 + \sin^2 \theta - 1$$

$$\Rightarrow 2 \left(\frac{\sin \theta}{\sin \theta} \right)^2 + \sin^2 \theta - 1$$

$$\Rightarrow 2 (1)^2 + \sin^2 \theta - 1$$

$$\Rightarrow 2 + \sin^2 \theta - 1$$

$$\Rightarrow \sin^2 \theta + 1$$

from given $\sin \theta = \cos \theta$

$$\text{so } \Rightarrow \theta = 45$$

$$\therefore \sin 45 = \cos 45$$

$$\frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}}$$

$$\therefore 2 \tan^2 \theta + \sin^2 \theta - 1 \Rightarrow \sin^2 \theta + 1$$

$$\Rightarrow \left(\frac{1}{\sqrt{2}} \right)^2 + 1$$

$$\Rightarrow \frac{1}{2} + 1$$

$$\Rightarrow \frac{3}{2}$$

$$\therefore 2 \tan^2 \theta + \sin^2 \theta - 1 = \underline{\underline{\frac{3}{2}}}$$

Solution - 26 (i)

$$\begin{aligned} \text{LHS} &\Rightarrow \cos \theta \cdot \tan \theta \\ &\rightarrow \cos \theta \cdot \frac{\sin \theta}{\cos \theta} \\ &\rightarrow \sin \theta \end{aligned}$$

$$\left(\because \tan \theta = \frac{\sin \theta}{\cos \theta} \right)$$

$$\text{RHS} = \sin \theta$$

$$\therefore \text{LHS} = \text{RHS}$$

Solution - 26 (ii)

$$\begin{aligned} \text{LHS} &\Rightarrow \sin \theta \cdot \cot \theta \\ &\rightarrow \sin \theta \cdot \frac{\cos \theta}{\sin \theta} \\ &\rightarrow \cos \theta \end{aligned}$$

$$\therefore \cot \theta = \frac{\cos \theta}{\sin \theta}$$

$$\therefore \text{LHS} = \text{RHS}$$

Solution - 26 (iii)

$$\begin{aligned} \text{LHS} &\Rightarrow \frac{\sin^2 \theta}{\cos \theta} + \cos^2 \theta \\ &\rightarrow \frac{\sin^2 \theta + \cos \theta \cdot \cos \theta}{\cos \theta} \\ &\rightarrow \frac{\sin^2 \theta + \cos^2 \theta}{\cos \theta} \\ &\rightarrow \frac{1}{\cos \theta} \end{aligned}$$


$$\left(\because \sin^2 \theta + \cos^2 \theta = 1 \right)$$

$$\therefore \text{LHS} = \text{RHS}$$

Solution- 27:

Given $\angle C = 90^\circ$

$$\tan A = \frac{3}{4}$$



By pythagorus theorem

$$AB^2 = BC^2 + AC^2$$

$$AB^2 = 3^2 + 4^2$$

$$AB^2 = 9 + 16$$

$$AB^2 = 25$$

$$AB = \sqrt{25}$$

$$AB = 5$$

$$\therefore \sin A = \frac{BC}{AB} = \frac{3}{5}$$

$$\cos A = \frac{AC}{AB} = \frac{4}{5}$$

$$\cos B = \frac{BC}{AB} = \frac{3}{5}$$

$$\sin B = \frac{AC}{AB} = \frac{4}{5}$$

$$\therefore \sin A \cos B + \cos A \sin B \Rightarrow \frac{3}{5} \cdot \frac{3}{5} + \frac{4}{5} \cdot \frac{4}{5}$$

$$\Rightarrow \frac{9}{25} + \frac{16}{25}$$

$$\Rightarrow \frac{9+16}{25}$$

$$\Rightarrow \frac{25}{25}$$

$$= 1$$

\therefore LHS = RHS.

Solution - 28 :- (a)

Given,

In $\triangle ABC$, $\triangle BRS$


$$AB = 18 \text{ cm}$$

$$BC = 7.5 \text{ cm}$$

$$RS = 5 \text{ cm}$$

$$\angle BSR = x^\circ$$

$$\angle SAB = y^\circ$$



From Hint ; $AR = 12 \text{ cm}$

$$RB = 6 \text{ cm}$$

$$AC = 19.5$$

$$(i) \tan x^\circ = \frac{RB}{SR} = \frac{6}{5}$$

$$(ii) \sin y = \frac{BC}{AC}$$

$$= \frac{7.5}{19.5} \times \frac{10}{10}$$

$$= \frac{75}{195}$$

~~3~~
13

$$= \frac{5}{13}$$

$$\therefore \sin y = \frac{5}{13}$$

Solution - 28 (b) :-

43

(i) From fig

By pythagorus theorem

$$AC^2 = AB^2 + BC^2$$


$$AC^2 = 12^2 + 5^2$$

$$AC^2 = 144 + 25$$

$$AC^2 = 169$$

$$AC = \sqrt{169}$$

$$AC = 13.$$



$$\begin{aligned} \therefore \cos \angle CBD &= \frac{AB}{AC} \\ &= \frac{12}{13} \end{aligned}$$

$$\begin{aligned} \text{(ii) } \cot \angle ABD &= \frac{BC}{AC} \\ &= \frac{5}{13} \end{aligned}$$

Solution - 24 :-


Given ABCD is rectangle

$$AC = 15$$

$$\angle ACD = \alpha ; \cot \alpha = \frac{3}{2}$$

$$\cot \alpha = \frac{CD}{AD} = \frac{3}{2}$$

$$\therefore CD = \frac{3}{2} AD$$



From $\triangle ACD$

$$AC^2 = AD^2 + CD^2$$

$$15^2 = AD^2 + \left(\frac{3}{2}AD\right)^2$$

$$225 = AD^2 + \frac{9AD^2}{4}$$

$$225 = \frac{4AD^2 + 9AD^2}{4}$$

$$225 \times 4 = 13AD^2$$

$$AD^2 = \frac{225 \times 4}{13}$$

$$AD = \sqrt{\frac{225 \times 4}{13}}$$

$$AD = \frac{15 \times 2}{\sqrt{13}}$$

$$AD = \frac{30}{\sqrt{13}}$$

$$\therefore CD = \frac{3}{2}AD$$

$$= \frac{3}{2} \cdot \frac{30}{\sqrt{13}}$$

$$= \frac{45}{\sqrt{13}}$$

\therefore Area of $\triangle ACD +$ Area of $\triangle ABC =$ Area of rectangle $ABCD$.

$$\therefore CD = \frac{45}{\sqrt{13}} = AB$$

$$AD = \frac{30}{\sqrt{13}} = BC$$

$$\therefore \text{Area of } \square ABCD = CD \times AD = \frac{45}{\sqrt{13}} \times \frac{30}{\sqrt{13}}$$

$$\begin{aligned}
 \text{Area} &= \frac{45 \times 30}{\sqrt{13} \cdot \sqrt{13}} \\
 &= \frac{1350}{13} \\
 &= 103 \frac{11}{13} \text{ cm}^2
 \end{aligned}$$

45

$$\begin{aligned}
 \therefore \text{Perimeter} &= 2(AB + BC) \\
 &= 2\left(\frac{45}{\sqrt{13}} + \frac{30}{\sqrt{13}}\right) \\
 &= 2\left(\frac{45 + 30}{\sqrt{13}}\right) \\
 &= \frac{2 \times 75}{\sqrt{13}} \\
 &= \frac{150}{\sqrt{13}}
 \end{aligned}$$

Solution - 30 :-

(a)

From $\triangle BCD$

\Rightarrow By Pythagoras theorem

$$BD^2 = BC^2 + CD^2$$


$$13^2 = 12^2 + CD^2$$

$$169 = 144 + CD^2$$

$$CD^2 = 169 - 144$$

$$CD^2 = 25$$

$$CD = \sqrt{25} = \underline{5}$$



$$(i) \sin \phi = \frac{CD}{BD}$$

$$= \frac{5}{13}$$

$$(ii) \tan \theta = \frac{DE}{AE}$$


$$= \frac{12}{9} = \frac{4}{3}$$

Solution - 30 (b) :-

$$\tan \theta = \frac{4}{3}$$

$$\therefore \sin \theta = \frac{DE}{AD}$$

$$AD = \frac{12}{\sin \theta}$$



$$(or) \cos \theta = \frac{AE}{AD}$$

$$AD = \frac{9}{\cos \theta}$$

Solution - 31 :- (1)

$$\text{LHS} \Rightarrow (\sin A + \cos A)^2 + (\sin A - \cos A)^2$$

$$\Rightarrow \sin^2 A + \cos^2 A + 2 \sin A \cos A + \sin^2 A + \cos^2 A - 2 \sin A \cos A$$

$$\left(\begin{array}{l} \therefore (a+b)^2 = a^2 + b^2 + 2ab \\ (a-b)^2 = a^2 + b^2 - 2ab \end{array} \right)$$

$$\Rightarrow 2 \sin^2 A + 2 \cos^2 A$$

$$\Rightarrow 2 (\sin^2 A + \cos^2 A)$$

$$\Rightarrow 2(1) = 2$$

$$\therefore \text{LHS} = \text{RHS}$$

Solution - 31 (ii) :-

$$\text{LHS} \Rightarrow \cot^2 A - \frac{1}{\sin^2 A} + 1$$

$$\Rightarrow \frac{\cos^2 A}{\sin^2 A} - \frac{1}{\sin^2 A} + 1$$

$$\Rightarrow \frac{\cos^2 A - 1 + \sin^2 A}{\sin^2 A}$$

$$\Rightarrow \frac{(\cos^2 A + \sin^2 A) - 1}{\sin^2 A}$$

$$\Rightarrow \frac{1 - 1}{\sin^2 A}$$

$$\Rightarrow 0 = \text{RHS}$$

Solution - 31 (iii) :-

$$\text{LHS} \Rightarrow \frac{1}{1 + \tan^2 A} + \frac{1}{1 + \cot^2 A}$$

$$\Rightarrow \frac{1}{1 + \frac{\sin^2 A}{\cos^2 A}} + \frac{1}{1 + \frac{\cos^2 A}{\sin^2 A}}$$

$$\Rightarrow \frac{1}{\frac{\cos^2 + \sin^2 A}{\cos^2 A}} + \frac{1}{\frac{\sin^2 A + \cos^2 A}{\sin^2 A}}$$

$$= \frac{\cos^2 A}{(1)} + \frac{\sin^2 A}{(1)}$$

$$\Rightarrow \cos^2 A + \sin^2 A$$

$$\Rightarrow 1 = \text{RHS} //$$

Solution - 32 :

48

$$\text{Given } \sqrt{\frac{1 - \sin^2 \theta}{1 - \cos^2 \theta}}$$

$$(\because \sin^2 \theta + \cos^2 \theta = 1)$$

$$\Rightarrow \sqrt{\frac{\cos^2 \theta}{\sin^2 \theta}}$$

$$\Rightarrow \sqrt{\cot^2 \theta}$$

$$\Rightarrow \cot \theta$$

Solution - 33 :-

$$\text{Given } \sin \theta + \operatorname{cosec} \theta = 2$$

Squaring on both sides.

$$(\sin \theta + \operatorname{cosec} \theta)^2 = 2^2$$

$$\sin^2 \theta + \operatorname{cosec}^2 \theta + 2 \sin \theta \cdot \operatorname{cosec} \theta = 4$$

$$\sin^2 \theta + \operatorname{cosec}^2 \theta + 2 \cancel{\sin \theta} \frac{1}{\cancel{\sin \theta}} = 4$$

$$\sin^2 \theta + \operatorname{cosec}^2 \theta = 4 - 2$$

$$\underline{\sin^2 \theta + \operatorname{cosec}^2 \theta = 2}$$

Solution - 34 :-

$$\text{Given } x = a \cos \theta + b \sin \theta$$

$$y = a \sin \theta - b \cos \theta$$

Squaring on b.s.

$$x^2 = (a \cos \theta + b \sin \theta)^2$$

$$x^2 = a^2 \cos^2 \theta + b^2 \sin^2 \theta + 2ab \sin \theta \cos \theta \quad \dots (1)$$

$$\therefore y = a \sin \theta - b \cos \theta$$

Squaring on b.s.

$$y^2 = (a \sin \theta - b \cos \theta)^2$$

$$y^2 = a^2 \sin^2 \theta + b^2 \cos^2 \theta - 2ab \sin \theta \cos \theta \quad \dots (2)$$

① + ②

$$\begin{aligned} \Rightarrow x^2 + y^2 &= a^2 \cos^2 \theta + b^2 \sin^2 \theta + 2ab \sin \theta \cos \theta + \\ &\quad a^2 \sin^2 \theta + b^2 \cos^2 \theta - 2ab \sin \theta \cos \theta \\ &= a^2 (\cos^2 \theta + \sin^2 \theta) + b^2 (\sin^2 \theta + \cos^2 \theta) \\ &= a^2 (1) + b^2 (1) \\ &= \underline{a^2 + b^2} \end{aligned}$$

\therefore LHS = RHS

Hence proved.